

Quantum discord and quantum phase transition in spin chains

Raoul Dillenschneider*

Department of Physics, University of Augsburg, D-86135 Augsburg, Germany

(Dated: December 17, 2008)

Quantum phase transitions of the transverse Ising and antiferromagnetic XXZ spin $S = 1/2$ chains are studied using *quantum discord*. Quantum discord allows the measure of quantum correlations present in many-body quantum systems. It is shown that the *amount* of quantum correlations increases close to the critical points. The observations are in agreement with the information provided by the concurrence which measures the entanglement of the many-body system.

PACS numbers: 03.67.-a, 64.70.Tg, 75.10.Pq

I. INTRODUCTION

Quantum phase transitions (QPT) occur when the ground state of a many-body system at absolute zero temperature undergoes a qualitative change by variation of a coupling and/or an external parameter¹. QPT are present in spin systems as for example in the one-dimensional XY model^{2,3} and in the antiferromagnetic XXZ model^{4,5,6}. Quantum phase transition in spin systems have been widely characterized considering the pairwise entanglement between two spin sites⁷. The concurrence which is a good measure of entanglement^{8,9} is maximal close to the critical points and its derivatives can signal more precisely the presence of a quantum phase transition at the critical points³.

In another register Ollivier and Zurek introduced the *quantum discord* as a measure of quantum correlations between two separated sub-systems of a many-body quantum system¹⁰. In information theory the *mutual information* is defined as the measure of correlation between two random variables $\mathcal{A}$ and $\mathcal{B}$. For classical system the mutual information can be expressed in terms of two equivalent expressions $\mathcal{I}(\mathcal{A}; \mathcal{B})$ and $\mathcal{J}(\mathcal{A}; \mathcal{B})$ (explicit definitions of each expressions of the mutual information are provided later in the text). However for quantum systems the two expressions are not equal and their difference gives the quantum discord. In parallel to the concurrence, quantum discord measures the quantum correlations of a quantum system. In contrary to the concurrence, the quantum discord can be different to zero even if a quantum system is separable. The Werner state, $\rho_{\text{Werner}} = \frac{1-\lambda}{4}\mathbb{I} + \lambda|\Psi\rangle\langle\Psi|$ where λ is a parameter and $|\Psi\rangle = (|00\rangle + |11\rangle)/\sqrt{2}$, provide an example for which the concurrence is equal to zero (separable state) for $\lambda < 1/3$ while the quantum discord is different to zero¹⁰. Hence quantum discord shows the existence of quantum correlations where concurrence shows that the system is separable. We point that the concurrence measures the *nonlocal quantumness* of correlations while quantum discord measures the total *amount* of quantum correlations of a state ρ .

Quantum discord is used in several context^{11,12} for its property to discernate quantum and classical correlations of a quantum systems. The behaviour of the quantum as

well as the classical correlations close to the critical points of QPTs can be studied. We focus our attention here on the quantum phase transitions of the one-dimensional Ising model which derives from the XY model and on the antiferromagnetic XXZ chain. Using the quantum discord we show that the *amount* of quantum correlations increases in the region close to the critical points. In opposite the *amount* of classical correlations close to the critical points decreases in the XXZ model while it is simply monotonous in the Ising model.

The outline of the paper is the following. In section II we recall the elements of information theory which leads to the construction of quantum discord as demonstrated by Ollivier and Zurek¹⁰. In section III and IV the quantum correlations close to the quantum phase transition of the Ising and XXZ models are studied. Appendix A provides the derivation of the quantum discord for spin $S = 1/2$ chains.

II. QUANTUM DISCORD

Information on the correlations between two random variables $\mathcal{A}$ and $\mathcal{B}$ can be obtained by measure of the *mutual information*. In classical information theory the mutual information reads

$$\mathcal{I}(\mathcal{A}; \mathcal{B}) = H(\mathcal{A}) + H(\mathcal{B}) - H(\mathcal{A}, \mathcal{B}), \quad (1)$$

where the information entropy $H(\mathcal{A}) = -\sum_a p_{\mathcal{A}=a} \log p_{\mathcal{A}=a}$ and $p_{\mathcal{A}=a}$ is the probability that a is the realisation of the random variable $\mathcal{A}$. The joint entropy of $\mathcal{A}$ and $\mathcal{B}$ is defined by $H(\mathcal{A}, \mathcal{B}) = -\sum_{a,b} p_{\mathcal{A}=a, \mathcal{B}=b} \log p_{\mathcal{A}=a, \mathcal{B}=b}$ where $p_{\mathcal{A}=a, \mathcal{B}=b}$ is the joint probability of the variables $\mathcal{A}$ and $\mathcal{B}$ to take respectively the values a and b . Using the Bayes rule $p_{\mathcal{A}, \mathcal{B}=b} = p_{\mathcal{B}=b} \times p_{\mathcal{A}|\mathcal{B}=b}$ the mutual information can be rewritten into the equivalent expression

$$\mathcal{J}(\mathcal{A}; \mathcal{B}) = H(\mathcal{A}) - H(\mathcal{A}|\mathcal{B}), \quad (2)$$

where $H(\mathcal{A}|\mathcal{B}) = -\sum_{a,b} p_{\mathcal{A}=a|\mathcal{B}=b} \log p_{\mathcal{A}=a|\mathcal{B}=b}$ is the conditional entropy of the random variables $\mathcal{A}$ and $\mathcal{B}$

and $p_{A=a|B=b}$ is the conditional probability of a being the realisation of the random variable $\mathcal{A}$ knowing that b is the realisation of the random variable $\mathcal{B}$. For classical random variables the two expressions (1) and (2) are strictly equivalent. However for quantum systems the two expressions of the mutual information are in general not equivalent and leads to define the *quantum discord* $\delta(\mathcal{A} : \mathcal{B}) = \mathcal{I}(\mathcal{A}; \mathcal{B}) - \mathcal{J}(\mathcal{A}; \mathcal{B})$ as demonstrated in refs.^{10,11}. Quantum discord is the difference between the two classically equivalent expressions of the mutual information.

The mutual information expressions (1) and (2) for quantum systems are obtained by replacing the Shannon entropy by the von Neumann entropy. The von Neumann entropy for the joint state reads $H(\mathcal{A}, \mathcal{B}) = -\text{Tr}_{\mathcal{A}, \mathcal{B}} \rho_{\mathcal{A}, \mathcal{B}} \ln \rho_{\mathcal{A}, \mathcal{B}}$ where the trace runs over the Hilbert spaces of the sub-systems $\mathcal{A}$ and $\mathcal{B}$, $\rho_{\mathcal{A}, \mathcal{B}}$ is the density matrix of the quantum system. In a similar way the entropy of the system $\mathcal{A}$ reads $H(\mathcal{A}) = -\text{Tr}_{\mathcal{A}} \rho_{\mathcal{A}} \log \rho_{\mathcal{A}}$ where $\rho_{\mathcal{A}}$ is the reduced density matrix obtained by taking the trace over all states of the system $\mathcal{B}$, $\rho_{\mathcal{A}} = \text{Tr}_{\mathcal{B}} \rho_{\mathcal{A}, \mathcal{B}}$. The quantum conditional entropy $H(\mathcal{A}|\mathcal{B})$ in equation (2) quantifies the ignorance of the state of $\mathcal{A}$ knowing the state of $\mathcal{B}$. It is worked out from the density matrix $\rho_{\mathcal{A}|\Pi_j^{\mathcal{B}}}$ which is related to the density matrix of the system $\rho_{\mathcal{A}, \mathcal{B}}$ through¹⁰

$$\rho_{\mathcal{A}|\Pi_j^{\mathcal{B}}} = \Pi_j^{\mathcal{B}} \rho_{\mathcal{A}, \mathcal{B}} \Pi_j^{\mathcal{B}} / p_j,$$

where $\Pi_j^{\mathcal{B}}$ is the projector onto the state j of the sub-system $\mathcal{B}$ and the probability $p_j = \text{Tr}_{\mathcal{A}, \mathcal{B}} \Pi_j^{\mathcal{B}} \rho_{\mathcal{A}, \mathcal{B}}$. The quantum conditional entropy then reads $H(\mathcal{A}|\mathcal{B}) = H(\mathcal{A} | \{\Pi_j^{\mathcal{B}}\}) = \sum_j p_j H(\rho_{\mathcal{A}|\Pi_j^{\mathcal{B}}})$.

Finally the quantum discord measuring the quantum correlations between the two quantum sub-systems $\mathcal{A}$ and $\mathcal{B}$ reads

$$\delta(\mathcal{A} : \mathcal{B}) = \min_{\{\Pi_j^{\mathcal{B}}\}} \left\{ H(\mathcal{A}) - H(\mathcal{A}, \mathcal{B}) + H(\mathcal{A} | \{\Pi_j^{\mathcal{B}}\}) \right\}.$$

It must be noticed that the quantum discord is minimized over the set of state $\{\Pi_j^{\mathcal{B}}\}$. The information obtained on the sub-system $\mathcal{A}$ is affected by the measure on the state of the sub-system $\mathcal{B}$. In order to get the maximum information on the sub-system $\mathcal{A}$ we need to consider the projection onto the states of $\mathcal{B}$ that disturb least the overall quantum system. Maximize the information on the sub-system $\mathcal{A}$ correspond to minimize the quantum discord with respects to the set of projector $\{\Pi_j^{\mathcal{B}}\}$.

Quantum discord provide information on the quantum nature of the correlations between two systems. If two sub-systems $\mathcal{A}$ and $\mathcal{B}$ are correlated *classically* the quantum discord is equal to zero. Moreover quantum discord shows that quantum correlations can be present for states that are not entangled. For example in Werner states the quantum correlations are still present while the system is separable¹⁰. To be more precise we can consider a quantum mixed state $\tilde{\rho} = |\psi\rangle\langle\psi|$ acting in an Hilbert space

$\mathcal{H}_{\mathcal{A}} \otimes \mathcal{H}_{\mathcal{B}}$. The quantum mixed state is separable if its entanglement is equal to zero. In this case the quantum separable state can be written as a product of quantum states and reads $\tilde{\rho} = \sum_i p_i |\psi_{i, \mathcal{A}}\rangle\langle\psi_{i, \mathcal{A}}| \otimes |\psi_{i, \mathcal{B}}\rangle\langle\psi_{i, \mathcal{B}}|$ where $|\psi_{i, \mathcal{A}}\rangle \in \mathcal{H}_{\mathcal{A}}$, $|\psi_{i, \mathcal{B}}\rangle \in \mathcal{H}_{\mathcal{B}}$ and $\sum_i p_i = 1$. The two systems $\mathcal{A}$ and $\mathcal{B}$ can be correlated through the separable state $\tilde{\rho}$ and the quantum states $|\psi_{i, \mathcal{A}}\rangle$ and $|\psi_{i, \mathcal{B}}\rangle$ do not have in general any classical counterpart. A consequence of the quantumness of the states $|\psi_{\mathcal{A}}\rangle$ and $|\psi_{\mathcal{B}}\rangle$ is that the correlations between the two systems $\mathcal{A}$ and $\mathcal{B}$ have quantum as well as classical nature. The quantum discord is non-zero while the entanglement of the quantum separable state is equal to zero. A good example is furnished by the two-qubit separable mixed state¹³

$$\tilde{\rho} = \frac{1}{4} \left(|\psi_+\rangle\langle\psi_+|_{\mathcal{A}} \otimes |0\rangle\langle 0|_{\mathcal{B}} + |\psi_-\rangle\langle\psi_-|_{\mathcal{A}} \otimes |1\rangle\langle 1|_{\mathcal{B}} + |0\rangle\langle 0|_{\mathcal{A}} \otimes |\psi_-\rangle\langle\psi_-|_{\mathcal{B}} + |1\rangle\langle 1|_{\mathcal{A}} \otimes |\psi_+\rangle\langle\psi_+|_{\mathcal{B}} \right), \quad (3)$$

where $|\psi_{\pm}\rangle = \frac{1}{\sqrt{2}} (|0\rangle \pm |1\rangle)$ and each qubit $\mathcal{A}$ and $\mathcal{B}$ have four nonorthogonal states. The two-qubit separable state (3) show a product of quantum states that can not have any equivalent classical system and also present quantum correlation. Quantum discord quantifies all quantum correlations including entanglement between $\mathcal{A}$ and $\mathcal{B}$.

Note that the mutual information $\mathcal{I}(\mathcal{A}; \mathcal{B})$ is the sum of the quantum discord and the classical correlation $CC(\mathcal{A}; \mathcal{B})$ defined in ref.¹⁴, $\mathcal{I}(\mathcal{A}; \mathcal{B}) = \delta(\mathcal{A} : \mathcal{B}) + CC(\mathcal{A}; \mathcal{B})$. In other words classical correlations $CC(\mathcal{A}; \mathcal{B})$ are equal to the mutual information $\mathcal{J}(\mathcal{A}; \mathcal{B})$.

Using quantum discord to discernate the quantum from the classical correlations we can study the behaviour of such different correlations in quantum phase transitions. In the next sections we focus our attention on the quantum phase transitions of the Ising and the XXZ spin-1/2 chains and we study them using the quantum discord.

III. QUANTUM DISCORD AND QUANTUM PHASE TRANSITION IN THE ISING CHAIN

The Hamiltonian of the transverse one-dimensionnal Ising model reads

$$H_{Ising} = - \sum_{i=1}^N (\lambda \sigma_i^x \sigma_{i+1}^x + \sigma_i^z),$$

with the boundary condition $\sigma_N^x = \sigma_1^x$ and σ^α with $\alpha = \{x, y, z\}$ are the Pauli matrices and $\sigma^0 = \mathbb{I}$. We denotes by $|g\rangle = |\uparrow\rangle$ and $|e\rangle = |\downarrow\rangle$ the spin up and spin down states. For $\lambda = 0$ all spins are pointing in

the z direction while for $\lambda \rightarrow \infty$ they point in the x direction. In the thermodynamic limit $N \rightarrow \infty$ the Ising spin chain undergoes a quantum phase transition at the critical point $\lambda_c = 1$. The correlation length diverges at this point^{6,7}.

The quantum discord is worked out from the joint state of the two spins at the lattice sites i and j . The information on the joint state is contained in the two-site density matrix ρ_{ij} which is derived from the following operator expansion

$$\rho_{ij} = \text{Tr}_{\bar{i}\bar{j}} [\rho] = \frac{1}{4} \sum_{\alpha, \beta=0}^3 \Theta_{\alpha\beta} \sigma_i^\alpha \otimes \sigma_j^\beta,$$

where the coefficients $\Theta_{\alpha\beta}$ of the expansion are related to the spin-spin correlation functions through the relation

$$\Theta_{\alpha\beta} = \text{Tr} [\sigma_i^\alpha \sigma_j^\beta \rho_{ij}] = \langle \sigma_i^\alpha \sigma_j^\beta \rangle.$$

Owing to the symmetry of the Hamiltonian most of the coefficients $\Theta_{\alpha\beta}$ are equal to zero. Translation invariance require that the density matrix verifies $\rho_{ij} = \rho_{i,i+r}$ and is independent of the position i . The reflexion symmetry leads to $\rho_{ij} = \rho_{ji}$, the Hamiltonian being real the density matrix verifies $\rho_{ij}^* = \rho_{ij}$ and the global rotation symmetry implies that the density matrix commutes with the operator $\sigma_i^z \sigma_j^z$. Combining all symmetry constraints the density matrix expressed in the natural basis $\{|gg\rangle, |ge\rangle, |eg\rangle, |ee\rangle\}$ reduces to⁶

$$\rho_{ij} = \begin{pmatrix} u_+ & 0 & 0 & y \\ 0 & w & x & 0 \\ 0 & x & w & 0 \\ y & 0 & 0 & u_- \end{pmatrix}, \quad (4)$$

with $u_{\pm} = \frac{1}{4} \pm \frac{\langle \sigma^z \rangle}{2} + \frac{\langle \sigma_i^z \sigma_j^z \rangle}{4}$, $w = \frac{1 - \langle \sigma_i^z \sigma_j^z \rangle}{4}$, $x = \frac{\langle \sigma_i^x \sigma_j^x \rangle + \langle \sigma_i^y \sigma_j^y \rangle}{4}$ and $y = \frac{\langle \sigma_i^x \sigma_j^x \rangle - \langle \sigma_i^y \sigma_j^y \rangle}{4}$. The magnetization of the spin-1/2 Ising chain is given by¹⁵

$$\langle \sigma^z \rangle = -\frac{1}{\pi} \int_0^\pi d\phi \frac{(1 + \lambda \cos \phi)}{\omega_\phi},$$

and $\omega_\phi = \sqrt{(\lambda \sin \phi)^2 + (1 + \lambda \cos \phi)^2}$ is the energy spectrum of the transverse Ising chain. The spin-spin correlations functions are related to the determinant of Toeplitz matrices¹⁶

$$\langle \sigma_i^x \sigma_{i+r}^x \rangle = \frac{\begin{vmatrix} G_{-1} & G_{-2} & \dots & G_{-r} \\ G_0 & G_{-1} & \dots & G_{-r+1} \\ \vdots & \vdots & \ddots & \vdots \\ G_{r-2} & G_{r-3} & \dots & G_{-1} \end{vmatrix}}{\begin{vmatrix} G_1 & G_0 & \dots & G_{-r+2} \\ G_2 & G_1 & \dots & G_{-r+3} \\ \vdots & \vdots & \ddots & \vdots \\ G_r & G_{r-1} & \dots & G_1 \end{vmatrix}},$$




FIG. 1: Concurrence for nearest neighbour spins in the Ising spin $S = 1/2$ chain. The concurrence is maximal close to the critical coupling $\lambda_c = 1$.

and $\langle \sigma_i^z \sigma_{i+r}^z \rangle = \langle \sigma^z \rangle^2 - G_r G_{-r}$ where

$$G_\kappa = \frac{1}{\pi} \int_0^\pi d\phi \cos(\phi\kappa) \frac{(1 + \lambda \cos \phi)}{\omega_\phi} - \frac{\lambda}{\pi} \int_0^\pi d\phi \sin(\phi\kappa) \frac{\sin \phi}{\omega_\phi}.$$

The quantum phase transition can be signaled by measure of entanglement. Indeed the nearest and next-nearest neighbours entanglement reach their maximum at the critical point λ_c ^{3,7}. A good measure of the entanglement is provided by the concurrence $\mathcal{C}$. The concurrence of two spins may be computed from the joint state ρ_{ij} through the formula $\mathcal{C} = \max\{0, \gamma_1 - \gamma_2 - \gamma_3 - \gamma_4\}$ where the γ_i are the eigenvalues in decreasing order of the matrix $R = \sqrt{\rho_{ij} \widetilde{\rho}_{ij}}$ ^{8,9}. The matrix $\widetilde{\rho}_{ij}$ is related to the transpose of the two-site density matrix by $\widetilde{\rho}_{ij} = (\sigma^y \otimes \sigma^y) \rho_{ij}^* (\sigma^y \otimes \sigma^y)$. Hence the entanglement between two spins at site i and j is given by

$$\mathcal{C}_{ij} = 2 \max\{0, |x| - \sqrt{u_+ u_-}, |y| - w\}.$$

Figures 1 and 2 represent respectively the concurrence for the nearest-neighbour spins $\mathcal{C}_{i,i+1}$ and next-nearest neighbour spins $\mathcal{C}_{i,i+2}$. It shows that the maximum of entanglement is reached close to $\lambda_c = 1$. Only the derivatives of the concurrence show singularities at the critical points³ signaling the presence of the quantum phase transition.

Entanglement increases close to the quantum phase transition. However entanglement is only one kind of quantum correlation. It is then legitimate to ask : What is the behaviour of the *total amount* of quantum correlations close to the critical point of the QPT ? The answer is provided by the quantum discord. The details of the derivation of the quantum discord are given in appendix A. Figure 3 represents the quantum discord for the Ising $S = 1/2$ chain for nearest-neighbour $\delta_{i,i+1}$ and




FIG. 2: Concurrence for next-nearest neighbour spins in the Ising spin chain. The concurrence is maximal at the critical coupling $\lambda_c = 1$.




FIG. 3: Quantum discord δ for the nearest-neighbour (in dashed line) and next-nearest-neighbour (in full line) spin sites of the one-dimensionnal Ising model. The quantum discord is minimized for $\phi = 0$ and θ varying progressively from zero for $\lambda < 1$ to $\pi/4$ for $\lambda > 1$, see appendix A for the details on the derivation.

next-nearest-neighbour $\delta_{i,i+2}$. It shows that the quantum correlations increase and are maximal in a region close to the critical point λ_c .

More informations on the location and the order of the quantum phase transition can be obtained by consideration of the derivatives of the quantum discord with respects to the coupling parameter λ . Indeed there is a relation between the nonanalyticity in energy and quantum discord that leads to the characterization of the quantum phase transition¹⁷. Consider the energy of the spin system in terms of the reduced density matrix of two spins at positions i and j , the energy reads

$$E(\rho_{ij}) = \sum_{ij} \text{Tr} [\tilde{H}_{ij} \rho_{ij}], \quad (5)$$

where $\tilde{H}_{ij}$ is the reduced Hamiltonian of the two spins at




FIG. 4: First derivatives of the quantum discord with respects to the coupling parameter λ for the nearest-neighbour (in dashed line) and next-nearest-neighbour (in full line) spin sites of the one-dimensionnal Ising model.

position i and j . The sum of $\tilde{H}_{ij}$ over the positions i, j is equal to the Hamiltonian of the spin system, $\sum_{ij} \tilde{H}_{ij} \equiv H$. For the one-dimensionnal Ising model the reduced Hamiltonian reads $\tilde{H}_{ij}^{Ising} = -(\lambda \sigma_i^x \sigma_j^x \tilde{\delta}_{i+1,j} + \sigma_i^z / N)$

where N is the number of spins and $\tilde{\delta}_{ij}$ is the Kronecker delta. The order as well as the location of a quantum phase transition can be characterized by the nonanalyticity of the energy. If the first derivative of the ground state energy presents a finite discontinuity then the quantum phase transition is of the first order. However if the first derivative is continuous while the second derivative shows discontinuity or divergence then the quantum phase transition is of the second order. Moreover the derivatives of the energy are related to the derivatives of the reduced density matrix and one can show that¹⁷

$$\begin{aligned} \frac{\partial E(\rho_{ij})}{\partial \lambda} &= \sum_{ij} \text{Tr} \left[\frac{\partial \tilde{H}_{ij}}{\partial \lambda} \rho_{ij} \right], \\ \frac{\partial^2 E(\rho_{ij})}{\partial \lambda^2} &= \sum_{ij} \left\{ \text{Tr} \left[\frac{\partial^2 \tilde{H}_{ij}}{\partial \lambda^2} \rho_{ij} \right] + \text{Tr} \left[\frac{\partial \tilde{H}_{ij}}{\partial \lambda} \frac{\partial \rho_{ij}}{\partial \lambda} \right] \right\}. \end{aligned}$$

Note that the derivatives of the reduced Hamiltonian are continuous with respects to the coupling parameter λ . Hence a discontinuity in the first derivative $\frac{\partial E(\rho_{ij})}{\partial \lambda}$ of the energy at the critical point implies a discontinuity at least of one of the reduced density matrices ρ_{ij} . Similarly a discontinuity or a singularity in the second derivatives of the energy $\frac{\partial^2 E(\rho_{ij})}{\partial \lambda^2}$ requires the divergence of at least one of the derivatives $\frac{\partial \rho_{ij}}{\partial \lambda}$ at the critical points. It becomes now evident that the quantum phase transition can be characterized by an analysis of the derivatives of the quantum discord with respects to the coupling parameter λ . The quantum discord being dependent on the reduced density matrix one deduces that i) a discontinuity in the quantum discord implies a discontinuity in ρ_{ij}




FIG. 5: Classical correlation CC for the nearest-neighbour (in dashed line) and next-nearest-neighbour (in full line) spin sites of the one-dimensionnal Ising model.

(hence the first derivative of the energy is discontinuous) and a first order QPT, ii) a singularity in the derivative of the quantum discord implies a discontinuity or a divergence of $\frac{\partial \rho_{ij}}{\partial \lambda}$ (the second derivative of the energy is discontinuous) and the QPT is of the second order.

Figure 4 shows that for nearest neighbours $\partial \delta_{i,i+1} / \partial \lambda$ presents a discontinuity at the critical point and for the next-nearest neighbours $\partial \delta_{i,i+2} / \partial \lambda$ shows a singularity. The fact that the quantum discord is continuous while its derivatives are discontinuous or singular at $\lambda = \lambda_c$ indicates that the precise location of the critical point is at $\lambda_c = 1$ and that the QPT is of the second order.

For $\lambda > 1$ and approaching the critical point from the right, the classical correlations decrease in opposite to the quantum discord which increases as depicted by the figures 3 and 5. For $\lambda < 1$ both classical and quantum correlations are decreasing when the coupling parameter λ decreases. Classical correlations monotonically increase with respects to λ while quantum discord show that the amplitude of the quantum correlations increase close to the critical point λ_c . From the mutual information we infer that the spins are uncorrelated for $\lambda \rightarrow 0$ as depicted in figure 6 and can be confirmed by the Hamiltonian which reduces to $H_{Ising, \lambda=0} = -\sum_{i=1}^N \sigma_j^z$. Close to the critical point the *amount* of quantum correlations is maximal as supported by the concurrence.

IV. QUANTUM DISCORD AND QUANTUM PHASE TRANSITION IN THE XXZ CHAIN

The hamiltonian of the one-dimensionnal XXZ spin-1/2 chain reads

$$H_{XXZ} = \sum_{\langle ij \rangle} (S_i^x S_j^x + S_i^y S_j^y + \Delta S_i^z S_j^z),$$

where the sum runs over the nearest-neighbour bonds




FIG. 6: Mutual information $\mathcal{I}(\mathcal{A}; \mathcal{B})$ for the nearest-neighbour (in dashed line) and next-nearest-neighbour (in full line) spin sites of the one-dimensionnal Ising model.

$\langle ij \rangle$. At the critical point $\Delta_c = 1$ the XXZ spin chain undergoes a quantum phase transition between an XY phase for $-1 < \Delta < 1$ and an Ising antiferromagnetic phase for $\Delta > 1$.

The density matrix of two nearest-neighbour spin sites is given by (4). However owing to the spin-flip symmetry the magnetization is equal to zero, $\langle \sigma^z \rangle = 0$ and the XXZ Hamiltonian is also symmetric with respects to the rotation in the xy plan which leads to the equality between the spin-spin correlation function in the xy plan $\langle \sigma_i^x \sigma_j^x \rangle = \langle \sigma_i^y \sigma_j^y \rangle$. Gathering all the symmetry constraints of the XXZ Hamiltonian the joint density matrix ρ_{ij} reduces to

$$\rho_{ij} = \begin{pmatrix} u & 0 & 0 & 0 \\ 0 & w & x & 0 \\ 0 & x & w & 0 \\ 0 & 0 & 0 & u \end{pmatrix},$$

with $u_{\pm} = u = \frac{1}{4} + \frac{\langle \sigma_i^z \sigma_j^z \rangle}{4}$, $w = \frac{1 - \langle \sigma_i^z \sigma_j^z \rangle}{4}$, $x = \frac{\langle \sigma_i^x \sigma_j^x \rangle}{2}$ and $y = 0$.

The spin-spin correlation functions between nearest-neighbour spin sites for $-1 < \Delta < 1$ are given by^{18,19,20}

$$\begin{aligned} \langle \sigma_i^x \sigma_{i+1}^x \rangle &= \frac{\cos \pi \Phi}{\pi^2} \int_{-\infty}^{\infty} \frac{dx}{\sinh x} \frac{x \cosh x}{\cosh^2(\Phi x)} \\ &\quad - \frac{1}{\pi \sinh(\pi \Phi)} \int_{-\infty}^{\infty} \frac{dx}{\sinh x} \frac{\sinh((1 - \Phi)x)}{\cosh \Phi x}, \end{aligned}$$

where $\Delta = \cos(\pi \Phi)$, and

$$\begin{aligned} \langle \sigma_i^z \sigma_{i+1}^z \rangle &= 1 - \frac{2}{\pi^2} \int_{-\infty}^{\infty} \frac{dx}{\sinh x} \frac{x \cosh x}{\cosh^2(\Phi x)} \\ &\quad + \frac{2 \cot(\pi \Phi)}{\pi} \int_{-\infty}^{\infty} \frac{dx}{\sinh x} \frac{\sinh((1 - \Phi)x)}{\cosh \Phi x}. \end{aligned}$$

And for $\Delta > 1$ the correlation function are given by^{21,22,23}




FIG. 7: Concurrence for nearest-neighbour spin sites in the XXZ spin $S = 1/2$ one-dimensionnal model.




FIG. 8: Quantum discord for nearest neighbour spins in the XXZ spin $S = 1/2$ one-dimensionnal model. The quantum discord is minimized for $\phi = 0$ and θ varying progressively from zero for $\lambda < 1$ to $\pi/4$ for $\lambda > 1$, see appendix A for the details on the derivation.

$$\langle \sigma_i^x \sigma_{i+1}^x \rangle = \int_{-\infty+i/2}^{\infty+i/2} \frac{dx}{\sinh(\pi x)} \times \left(\frac{x}{\sin^2(\phi x)} \cosh \nu - \frac{\cot(\nu x)}{\sinh \nu} \right),$$

with $\Delta = \cosh \nu$, and

$$\langle \sigma_i^z \sigma_{i+1}^z \rangle = 1 + 2 \int_{-\infty+i/2}^{\infty+i/2} \frac{dx}{\sinh(\pi x)} \times \left(\cot(\nu x) \coth(\nu) - \frac{x}{\sin^2(\nu x)} \right).$$

The concurrence for the XXZ spin-1/2 chain is given by $\mathcal{C} = 2 \max \{0, |x| - |u|\}$. Figure 7 represents the dependence of the concurrence with respects to the control parameter Δ and shows that entanglement between the




FIG. 9: Dependence of the classical correlation $CC(\mathcal{A}; \mathcal{B})$ on the parameter Δ for the nearest-neighbour spin sites in the XXZ spin $S = 1/2$ chain.

nearest-neighbour spin sites is maximal at the critical point $\Delta_c = 1$ ⁶.

In parallel the quantum discord presents the maximum of the quantum correlations at the critical point. The derivation of the quantum discord for the XXZ model follow the same scheme as for the Ising model and the details of the derivation are provided in appendix A. Figure 8 depicts the dependence of the quantum discord on the control parameter Δ . The quantum discord is minimized with respects to the angles ϕ and θ that control the axes of the projector $\Pi_j^{\mathcal{B}}$.

As depicted in figure 8 the quantum correlations are stronger close to the QPT. On the opposite the classical correlations are weaker close to the critical point Δ_c . Both classical and quantum correlations measured by quantum discord and the mutual information $\mathcal{I}(\mathcal{A}; \mathcal{B})$ behave accordingly to the behaviour of entanglement measured by the concurrence.

The cusps observed in figures 8 and 9 for both the classical correlation and the quantum discord arise from a level-crossing between the ground-state energy and an excited-state energy. This scheme explains the cusp observed in the concurrence measuring the entanglement of nearest neighbour spin in $(d \geq 2)$ -dimensionnal XXZ model⁴. A similar mechanism explains also the fact that the concurrence is maximal at the critical points for the one-dimensionnal XXZ model²⁴. Level crossing induces a qualitative change in the ground states at the critical point $\Delta_c = 1$ and leads to the abrupt behaviour of the quantum discord and classical correlation at the quantum phase transition. As depicted in figure 7 the concurrence is less sensitive to the level-crossing than the quantum discord and the classical correlations since it does not present an abrupt cusp at Δ_c .

Remark that the quantum discord for the next-nearest neighbour spins $\delta_{i,i+2}$ is not presented for the 1d XXZ model. The reason is that no more informations can be gained by $\delta_{i,i+2}$ than already furnished by the quantum

discord for nearest neighbours $\delta_{i,i+1}$. Indeed the quantum discord $\delta_{i,i+2}$ presents a similar dependence on the coupling parameter Δ as shown by $\delta_{i,i+1}$ and its amplitude verifies $\delta_{i,i+1} \sim 2\delta_{i,i+2}$.

V. CONCLUSION

Quantum phase transition of the one-dimensional spin-1/2 Ising and antiferromagnetic XXZ models have been studied using the quantum discord^{10,11}. Quantum discord is a measure of the quantum correlation given by the difference of two classically equivalent expressions of the mutual information. Using quantum discord and mutual information we showed that it is possible to discernate the quantum and the classical nature of the correlations between two elements of a quantum many-body system.

We demonstrated that the amount of quantum correlations is larger close to the critical points for the quantum phase transitions of the one-dimensionnal XXZ and Ising models. For the XXZ model the amount of classical correlations decrease at the critical point while for the Ising model they are simply monotonous with respects to the coupling parameter λ . The behaviours of the quantum and classical correlations measured by means of the quantum discord are in agreement with the behaviour of the entanglement computed by the concurrence for both spin models.

Quantum discord is a good way to discernate the nature of the correlations between the components of a quantum system and it is a good qualitative indicator of the existence of quantum phase transition. However depending on the model studied the maximum of the quantum discord is not necessarily located at the critical points as shown for the Ising model. The quantum discord can only be used for qualitative detection of quantum phase transitions. On the contrary the derivatives of the quantum discord provide precise informations on the location and on the order of the quantum phase transitions. Quantum discord^{10,11} and concurrence^{4,5,6,7} are two useful tools to detect quantum phase transitions. Despite the fact that implementation of concurrence is more easy/rapid than for quantum discord, concurrence measures only the nonlocal quantumness of the correlations between two systems $\mathcal{A}$ and $\mathcal{B}$. Quantum discord measures the total amount of quantum correlations between $\mathcal{A}$ and $\mathcal{B}$ including entanglement.

We showed that quantum discord agrees with the predictions provided by the concurrence for the behaviour of the quantum correlations close to quantum phase transitions for one-dimensionnal quantum systems. It would be interesting to confirm this observation in futur investigation of quantum systems with higher dimensionnality.

Acknowledgments

The author would like to thank Eric Lutz for useful

discussions. This work was supported by the Emmy Noether Program of the DFG (Contract LU1382/1-1) and the cluster of excellence Nanosystems Initiative Munich (NIM).

APPENDIX A: DERIVATION OF THE QUANTUM DISCORD FOR SPIN CHAINS

The quantum discord for two systems reads $\delta(\mathcal{A} : \mathcal{B}) = \min_{\{\Pi_j^{\mathcal{B}}\}} \left\{ H(\mathcal{A}) - H(\mathcal{A}, \mathcal{B}) + H(\mathcal{A} | \{\Pi_j^{\mathcal{B}}\}) \right\}$ for which the three von Neumann entropy are computed over respectively the reduced density matrix $\rho_{\mathcal{A}}$, the joint state $\rho_{\mathcal{A},\mathcal{B}}$ and the conditional density matrix $\rho_{\mathcal{A}|\Pi_j^{\mathcal{B}}}$.

The reduced density matrix $\rho_{\mathcal{A}}$ is given by

$$\begin{aligned} \rho_{\mathcal{A}} &= \text{Tr}_{\mathcal{B}} \rho_{\mathcal{A},\mathcal{B}} \\ &= (u_+ + w) |g\rangle\langle g|_{\mathcal{A}} + (u_- + w) |e\rangle\langle e|_{\mathcal{A}}, \end{aligned}$$

which leads to the von Neumann entropy of the reduced density matrix

$$\begin{aligned} H(\mathcal{A}) &= -\text{Tr}_{\mathcal{A}} \rho_{\mathcal{A}} \log \rho_{\mathcal{A}} \\ &= -\frac{1}{2} (1 + \langle \sigma^z \rangle) \log \left[\frac{1}{2} (1 + \langle \sigma^z \rangle) \right] \\ &\quad - \frac{1}{2} (1 - \langle \sigma^z \rangle) \log \left[\frac{1}{2} (1 - \langle \sigma^z \rangle) \right]. \quad (\text{A1}) \end{aligned}$$

The joint density matrix is equal to the density matrix given in equation (4) $\rho_{\mathcal{A},\mathcal{B}} = \rho_{ij}$ and the von Neumann entropy is given yb the sum of the Shannon entropy over the eigenvalues of the density matrix ρ_{ij} and reads

$$\begin{aligned} H(\mathcal{A}, \mathcal{B}) &= -\text{Tr}_{\mathcal{A},\mathcal{B}} \rho_{\mathcal{A},\mathcal{B}} \log \rho_{\mathcal{A},\mathcal{B}} \\ &= -(w+x) \log(w+x) - (w-x) \log(w-x) \\ &\quad - \sum_{\epsilon=\pm} \Xi_{\epsilon} \log \Xi_{\epsilon}, \quad (\text{A2}) \end{aligned}$$

where

$$\Xi_{\pm} = \frac{1}{2} \left\{ (u_+ + u_-) \pm [(u_+ - u_-)^2 + 4y^2]^{1/2} \right\}.$$

The conditional density matrix $\rho_{\mathcal{A}|\Pi_j^{\mathcal{B}}}$ is given by projection on the arbitrary basis

$$\begin{aligned} |g\rangle_{\mathcal{B}} &= \cos(\theta) |j_1\rangle_{\mathcal{B}} + e^{i\phi} \sin(\theta) |j_2\rangle_{\mathcal{B}}, \\ |e\rangle_{\mathcal{B}} &= e^{-i\phi} \sin(\theta) |j_1\rangle_{\mathcal{B}} - \cos(\theta) |j_2\rangle_{\mathcal{B}}. \end{aligned}$$

The angles ϕ and θ control the projectors direction and the quantum discord is minimized with respects to these angles.

The conditional entropy is then given by

$$\rho_{\mathcal{A}|\Pi_j^{\mathcal{B}}} = \Pi_j^{\mathcal{B}} \rho_{\mathcal{A},\mathcal{B}} \Pi_j^{\mathcal{B}} / p_j,$$

with $\Pi_{j\alpha=1,2}^{\mathcal{B}} = \mathbb{I}_{\mathcal{A}} \otimes |j_{\alpha}\rangle\langle j_{\alpha}|_{\mathcal{B}}$ and $p_{j\alpha} = \text{Tr}_{\mathcal{A},\mathcal{B}} \mathbb{I}_{\mathcal{A}} \otimes |j_{\alpha}\rangle\langle j_{\alpha}|_{\mathcal{B}} \rho_{\mathcal{A},\mathcal{B}}$.

For the projection onto the state $|j_\alpha\rangle$ where $\alpha = 1, 2$ one gets

$$\rho_{\mathcal{A}|\Pi_{j_\alpha}^{\mathcal{B}}} = |j_\alpha\rangle\langle j_\alpha|_{\mathcal{B}} \otimes \left\{ |g\rangle\langle g|_{\mathcal{A}} X_{j_\alpha,+} + |e\rangle\langle e|_{\mathcal{A}} X_{j_\alpha,-} + |g\rangle\langle e|_{\mathcal{A}} Y_{j_\alpha} + |e\rangle\langle g|_{\mathcal{A}} \bar{Y}_{j_\alpha} \right\} / p_{j_\alpha}, \quad (\text{A3})$$

where for $\alpha = 1$,

$$\begin{aligned} X_{j_1,+} &= (u_+ \cos^2(\theta) + w \sin^2(\theta)), \\ X_{j_1,-} &= (u_- \sin^2(\theta) + w \cos^2(\theta)), \\ Y_{j_1} &= (e^{-i\phi} x + y e^{i\phi}) \sin(\theta) \cos(\theta), \end{aligned}$$

and with $p_{j_1} = \frac{1}{2}(1 + \langle \sigma^z \rangle \cos(2\theta))$. For the projection onto the state $|j_2\rangle$ the elements $X_{j_2,\pm}$ are given by

$$\begin{aligned} X_{j_2,+} &= (u_+ \sin^2(\theta) + w \cos^2(\theta)), \\ X_{j_2,-} &= (u_- \cos^2(\theta) + w \sin^2(\theta)), \\ Y_{j_2} &= -(e^{-i\phi} x + y e^{i\phi}) \sin(\theta) \cos(\theta), \end{aligned}$$

and with $p_{j_2} = \frac{1}{2}(1 - \langle \sigma^z \rangle \cos(2\theta))$. The eigenvalues $\Psi_{j_\alpha,\pm}$ of the conditional density matrix

$$(\text{A3}) \quad \text{reads} \quad \Psi_{j_\alpha,\pm} = \frac{1}{2p_{j_\alpha}} \left\{ (X_{j_\alpha,+} + X_{j_\alpha,-}) \pm [(X_{j_\alpha,+} - X_{j_\alpha,-})^2 + 4Y_{j_\alpha} \bar{Y}_{j_\alpha}]^{1/2} \right\}.$$

Finally the conditional von Neumann entropy reads

$$\begin{aligned} H(\mathcal{A}|\{\Pi_j^{\mathcal{B}}\}) &= \sum_{j=1,2} -p_j \text{Tr}_{\mathcal{A}} \rho_{\mathcal{A}|\Pi_j^{\mathcal{B}}} \log \rho_{\mathcal{A}|\Pi_j^{\mathcal{B}}} \\ &= - \sum_{\epsilon=\pm} \sum_{j=1,2} p_j \Psi_{j,\epsilon}(\phi, \theta) \log \Psi_{j,\epsilon}(\phi, \theta). \end{aligned} \quad (\text{A4})$$

The quantum discord is minimized with respects to the angles ϕ and θ that control the direction of the projector $\Pi_j^{\mathcal{B}}$ in the Hilbert space of the sub-system $\mathcal{B}$. The minimization must be achieved for each ensemble of values of the magnetization and the spin-spin correlations function which defines the joint density matrix. It can be shown numerically that for the one-dimensionnal Ising and antiferromagnetic XXZ models the set of angles $\{\phi, \theta\}$ that minimize the quantum discord belong to the ranges $\phi = 0, \theta \in [0, \pi/4]$.

-
- * E-mail address : raoul.dillenschneider@physik.uni-augsburg.de
- ¹ S. Sachdev, *Quantum phase transitions* (Cambridge Univ. Press, Cambridge, 2000).
 - ² L. Amico, R. Fazio, A. Osterloh, V. Vedral, Rev. Mod. Phys. **80**, 517-576 (2008).
 - ³ A. Osterloh, L. Amico, G. Falci, R. Fazio, Nature **416**, 608 (2002).
 - ⁴ S.-J. Gu, G.-S. Tian, H.-Q. Lin, Phys. Rev. A **71**, 052322 (2005).
 - ⁵ S.-J. Gu, G.-S. Tian, H.-Q. Lin, New Journal of Physics **8**, 61 (2006).
 - ⁶ J.-M. Cai, Z.-W. Zhou, G.-C. Guo, Phys. Lett. A **352**, 196 (2006).
 - ⁷ T. J. Osborne, M. A. Nielsen, Phys. Rev. A **66**, 032110 (2002).
 - ⁸ W. K. Wothers, Phys. Rev. Lett. **80**, 2245 (1998).
 - ⁹ K. M. O'Connor, W. K. Wothers, Phys. Rev. A **63**, 052302 (2001).
 - ¹⁰ H. Ollivier and H. Zurek, Phys. Rev. Lett. **88**, 017901 (2002).
 - ¹¹ H. Zurek, Phys. Rev. A **67**, 012320 (2003).
 - ¹² R. Dillenschneider, E. Lutz, arXiv:0803.4067 (2008).
 - ¹³ A. Datta, A. Shaji, C.M. Caves, Phys. Rev. Lett. **100**, 050502 (2008).
 - ¹⁴ L. Henderson, V. Vedral, J. Phys. A : Math. Gen. **34**, 6899 (2001).
 - ¹⁵ E. Barouch, B.M. McCoy, Phys. Rev. A **2**, 1075 (1970).
 - ¹⁶ E. Barouch, B.M. McCoy, Phys. Rev. A **3**, 786 (1971).
 - ¹⁷ L.-A. Wu, M.S. Sarandy, D.A. Lidar, Phys. Rev. Lett. **93**, 250404 (2004).
 - ¹⁸ M. Jimbo, T. Miwa, J. Phys. A **29**, 2923 (1996).
 - ¹⁹ V.E. Korepin, A.G. Izergin, F.H.L. Essler, D.B. Uglov, Phys. Lett. A **190**, 182 (1994).
 - ²⁰ G. Kato, M. Shiroishi, M. Takahashi, K. Sakai, J. Phys. A **36**, L337 (2003).
 - ²¹ M. Jimbo, K. Miki, T. Miwa, A. Nakayashiki, Phys. Lett. A **168**, 256 (1992).
 - ²² A. Nakayashiki, Int. J. Mod. Phys. A **9**, 5673 (1994).
 - ²³ M. Takahashi, G. Kato, M. Shiroishi, J. Phys. Soc. Japan **73**, 245 (2004).
 - ²⁴ S.-J. Gu, G.-S. Tian, H.-Q. Lin, Chinese Phys. Lett. **24**, 2737 (2007).